

ON THE FIFTH ATOM-BOND CONNECTIVITY INDEX OF ARMCHAIR POLYHEX NANOTUBE

K. Pattabiraman¹, P. Kandan¹

¹Department of Mathematics, Faculty of Engineering and Technology
Annamalai University, Annamalainagar, 608 002, India
e-mail: pramank@gmail.com

Abstract. Among topological descriptors, connectivity indices are very important and they have a prominent role in chemistry. One of them is atom-bond connectivity (ABC) index of a graph G and defined as $ABC(G) = \sum_{e=uv \in E(G)} \sqrt{\frac{(Q_u + Q_v - 2)}{Q_u Q_v}}$, where Q_v is some quantity that in a unique manner can be associated with the vertex u of graph G . Recently, Calimi introduced a new version of atom-bond connectivity (ABC_5) index as $ABC_5(G) = \sum_{e=uv \in E(G)} \sqrt{\frac{(M_u + M_v - 2)}{M_u M_v}}$, where M_u denotes the products of the degrees of adjacent vertices of u . In this paper, formula for calculating the above topological descriptor in armchair polyhex nanotube $TUAC_6[m, n]$ $m, n \geq 1$ family is given.

Keywords: nanotubes, fourth atom-bond connectivity index, armchair polyhex.

AMS Subject Classification: 05C12, 05C76.

1. Introduction

All the graphs considered in this paper are simple and connected. Graph theory has successfully provided chemists with a variety of useful tools [3, 4, 5, 6], among which are the topological indices. In theoretical chemistry, assigning a numerical value to the molecular structure that will closely correlate with the physical quantities and activities. Molecular structure descriptors (also called topological indices) are used for modeling physicochemical, pharmacologic, toxicologic, biological and other properties of chemical compounds. Let $G = (V, E)$ be a simple molecular graph without directed and multiple edges and without loops, the vertex and edge sets of it are represented by $V = V(G)$ and $E = E(G)$, respectively. In chemical graphs, the vertices correspond to the atoms of the molecule, and the edges represent to the chemical bonds. Also, if e is an edge of G , connecting the vertices u and v , then we write $e = uv$ and say u and v are adjacent. A connected graph is a graph such that there is a path between all pairs of vertices. Among topological descriptors, connectivity indices are very important and they have a prominent role in chemistry. One of the best known and widely used is the connectivity index, introduced in 1975 by Milan Randić [7], who has shown this index to reflect molecular branching and defined as follows:

$\chi(G) = \sum_{e=uv \in E(G)} \frac{1}{\sqrt{d_u d_v}}$, where d_u denotes the degree of vertex u of G . Another important classes of connectivity indices is *atom-bond connectivity (ABC) index*, introduced by Furtula et al.[8], by setting Q_u and Q_v to be the degree of a vertex u and v , defined as $ABC_1(G) = \sum_{e=uv \in E(G)} \sqrt{\frac{(d_u+d_v-2)}{d_u d_v}}$, which it has been applied up until now to study the stability of alkanes and the strain energy of cycloalkanes. The second and third members of this class were introduced by A. Graovac and M. Ghorbani[10] and M. R. Farahani,[9] on distance version, separately. The fourth member of this class was considered by M. Ghorbani et al.[11], by setting Q_u to be the sum of degrees of all vertices adjacent to vertex u . In this extend, Calimli[12] proposed the new atom bond connectivity (ABC_5) index and obtained the exact value for infinite class of dendrimers, defined as follows $ABC_5(G) = \sum_{e=uv \in E(G)} \sqrt{\frac{(M_u+M_v-2)}{M_u M_v}}$, where M_u denotes the products of the degrees of adjacent vertices of u . On the continuation of this work, we consider another infinite class molecular graphs armchair polyhex nanotube $TUAC_6[m, n]$. In Refs [13, 14, 15, 16, 17, 18, 19] some topological indices of armchair polyhex nanotube $TUAC_6[m, n]$ (Figure 1) is computed. In this paper, we continue this work to compute the fifth atom-bond connectivity index of molecular graphs related to armchair polyhex nanotube $TUAC_6[m, n]$

2. Result

The aim of this section is to obtain a closed formula of ABC_5 of general representation of armchair polyhex nanotubes $TUAC_6[m, n]$



Figure 1. The 3D Lattice of Armchair polyhex nanotubes $TUAC_6[8,7]$.

Theorem 2.1. Let G be the armchair nanotube $TUAC_6[m, n]$, for every $m, n \geq 1$. Then the fifth atom

bond connectivity index of G is equal to $(\frac{\sqrt{10}}{6} + \frac{2}{3}\sqrt{\frac{11}{6}} + \frac{1}{9}\sqrt{\frac{17}{2}} + \frac{2}{9}\sqrt{\frac{43}{6}} + \frac{2n}{3}\sqrt{13} - \frac{8}{27}\sqrt{13})m$.

Proof. Consider the armchair polyhex nanotubes $G = TUAC_6[m, n]$ (Figure 1). Suppose m and n denote the number of hexagons in the first row/column of the 2D- lattice of $TUAC_6[m, n]$ (Figure 2), respectively. Thus the number of vertices/atoms in this nanotube is equal to $|V(G)| = 2m(n + 1)$, $m, n \geq 1$ and obviously the number of edges/bonds is $|E(G)| = 3mn + 2m$. For the convinence let us partition the vertex set and edge set of G as follows. Since the degree of an arbitrary vertex/atom of a molecular graph armchair polyhex is equal to 2. Thus let $V(G) = V_1 \cup V_2$, where $V_1 = \{v \in V(G) | d_v = 2\}$ and $V_2 = \{v \in V(G) | d_v = 3\}$ and let $E(G) = E_1 \cup E_2 \cup E_3$, where $E_1 = \{uv \in E(G) | d_u = d_v = 3\}$, $E_2 = \{uv \in E(G) | d_u = 3, d_v = 2\}$ and $E_3 = \{uv \in E(G) | d_u = d_v = 2\}$. From Figure 2, it is easy to see that the size of edge/bond partitions E_3, E_2 and E_1 are equal to $m, 2m$ and $3mn - m$, respectively and one can observe that for every atom/vertex $v \in V_1, M_v = 2 \times 3 = 6$, since for its adjacent vertices u, w , observe that $d_u = 2$ and $d_w = 3$ ($u \in V_1, w \in V_2$) and obviously $M_u = 2 \times 3 = 6$, whereas $M_w = 3 \times 3 \times 2 = 18$, since for $N_w = \{u_1, u_2, v\}$, the degree of vertices/atoms u_1, u_2 , is equal to three. Also, for all other vertices x (which belong to V_2), $S_x = 3 \times 3 \times 3 = 27$.




Figure 2. The 2D Lattice of Armchair polyhex nanotubes $TUAC_6[8,7]$.

From Figure 2, the first and end ($1^{th}, 2n + 1^{th}$) rows consider all members of V_1 and $|V_1| = 2m$, also all similar vertices to w belong to second and $2n^{th}$ rows. Therefore $\forall m, n \geq 1$, we have following computations for the fifth atom-bond connectivity index of armchair polyhex $G = TUAC_6[m, n]$:

$$ABC_5(G) = \sum_{e=uv \in E(G)} \sqrt{\frac{(M_u + M_v - 2)}{M_u M_v}}$$

$$\begin{aligned}
&= \sum_{e=uv \in E_1} \sqrt{\frac{(M_u+M_v-2)}{M_u M_v}} + \sum_{e=uv \in E_2} \sqrt{\frac{(M_u+M_v-2)}{M_u M_v}} + \\
&\sum_{e=uv \in E_3} \sqrt{\frac{(M_u+M_v-2)}{M_u M_v}} \\
&= m \sqrt{\frac{6+6-2}{6 \times 6}} + 2m \sqrt{\frac{6+18-2}{6 \times 18}} + m \sqrt{\frac{18+18-2}{18 \times 18}} + 2m \sqrt{\frac{18+27-2}{18 \times 27}} + \\
&(9mn - 4m) \sqrt{\frac{27+27-2}{27 \times 27}} \\
&= \frac{m}{6} \sqrt{10} + \frac{2m}{3} \sqrt{\frac{11}{6}} + \frac{m}{9} \sqrt{\frac{17}{2}} + \frac{2m}{9} \sqrt{\frac{43}{6}} + \frac{2(9mn-4m)}{27} \sqrt{13} \\
&= \left(\frac{\sqrt{10}}{6} + \frac{2}{3} \sqrt{\frac{11}{6}} + \frac{1}{9} \sqrt{\frac{17}{2}} + \frac{2}{9} \sqrt{\frac{43}{6}} + \frac{2n}{3} \sqrt{13} - \frac{8}{27} \sqrt{13} \right) m \\
&\approx (2.40730n + 1.28025) m
\end{aligned}$$

Example 2.2. For $TUAC_6[8,7]$ with 128 atoms and 184 chemical bonds. Then one can see that

$$ABC_5(TUAC_6[8,7]) = 144.8492.$$

References

1. Klein D.J., Doslić T., Bonchev D., Vertex-weightings for distance moments and thorny graphs, *Discrete Appl. Math.*, Vol.155, No.17, 2007, pp.2294-2302.
2. Devillers J., Balaban A.T., Eds., *Topological indices and related descriptors in QSAR and QSPR*, Gordon and Breach, Amsterdam, The Netherlands, 1999.
3. Balaban A.T. (Ed.), *Chemical Applications of Graph Theory*, Academic Press, London, 1976.
4. Graovac A., Gutman I., Vukićević D. (Ed.), *Mathematical Methods and Modelling for Students of Chemistry and Biology*, Hum, Zagreb, 2009.
5. Gutman I., *Introduction to Chemical Graph Theory*, Fac. Sci. Kragujevac, Kragujevac, 2003 (in Serbian).
6. Gutman I., (Ed.), *Mathematical Methods in Chemistry*, Prijepolje Museum, Prijepolje, 2006.
7. Randić M., On characterization of molecular branching, *J. Amer. Chem. Soc.*, 97, 1975, pp.6609-6615.
8. Furtula B., A. Graovac, Vukićević D., Atom-bond connectivity index of trees, *Discrete Appl. Math.*, 157, 2009, pp.2828-2835.

9. Farahani M.R., A New Version of Atom-Bond Connectivity Index of Circumcoronene Series of Benzenoid, *Pacific Journal of Applied Mathematics*, Vol.5, No.4, 2013, pp.245-252.
10. Graovac A., Ghorbani M., A New Version of Atom-Bond Connectivity Index. *Acta Chim. Slov.*, 57, 2010, pp.609-612.
11. Ghorbani M., Hosseinzadeh M.A., Computing ABC4 index of nanostar dendrimers, *Optoelectron, Adv. Mater.-Rapid Commun.*, Vol.4, No.9, 2010, pp.1419-1422.
12. Calimli M.H., The Fifth Version of Atom Bond Connectivity Index (ABC5) of an infinite class of dendrimers, *Optoelectron. Adv. Mater.-Rapid Commun.*, Vol.5, No.10, 2011, pp.1091-1092.
13. Ashrafi A.R., Loghman A., PI index of armchair polyhex nanotubes, *ARS Combinatoria*, 80, 2006, pp.193-199.
14. Iranmanesh A., Ashrafi A.R., Balaban index of an armchair polyhex TUC4C8(R) and TUC4C8(S) nanotorus, *Journal of Computational and Theoretical Nanoscience.*, Vol.4, No.3, 2007, pp.514-517.
15. Deng H., The PI Index of TUAC6[2p,q], *MATCH Communications in Mathematical and in Computer Chemistry*, 55, 2006, pp.461-476.
16. Diudea M.V., Stefu M., P'arv B., John P.E., Armchair Polyhex Nanotubes, *Croatica Chemica Acta.*, 77, 2004, pp.111-115.
17. Eliasi M., Taeri B., Distance in Armchair Polyhex Nanotubes, *MATCH Comms. in Math. and in Comp. Chem.*, 62, 2009, pp.295-310.
18. Yousefi S., Ashrafi A.R., Distance Matrix and Wiener Index of Armchair Polyhex Nanotubes, *Studia Univ. Babeş-Bolyai. Chemia*, Vol.53, No.4, 2008, pp.111-116.
19. Farahani M.R., On the Fourth Atom Bond Connectivity Index of Armchair Polyhex Nanotubes, *Proc. Rom. Acad., Series B*, Vol.15, No.1, 2013, pp.3-6.