

PROSOPIS AFRICANA EXTRACTS AS POTENTIAL NATURAL ALTERNATIVES TO SYNTHETIC ANTIBIOTICS AND A KEY FOR SUSTAINABLE BROILER PRODUCTION: A REVIEW

Alagbe Olujimi John*

Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India

Abstract. The chicken sector is more vulnerable to antimicrobial resistance, the buildup of toxic or dangerous residues in meat and eggs and environmental contamination as a result of the widespread and careless use of antibiotics in most nations. Natural solutions, such as medicinal plants, are required to address these issues. These plants have been shown to contain a variety of phytochemicals, including phenols, flavonoids, terpenoids, tannins, alkaloids and saponins. These chemicals endow plants with a multitude of medicinal qualities. One of the many alternatives to antibiotics is *Prosopis africana* extract, which includes stem bark, leaves, roots and their essential oils. Antioxidant, antibacterial, antifungal, anti-helminthic, antiviral, hepatoprotective, immune-stimulatory and antimicrobial properties are among the many pharmacological activity of *P. africana* preparations. They are also an abundant supplier of vital minerals and amino acids that support enzyme function and provide defense against the effects of oxidative stress. Using extracts from *Prosopis africana* is the way forward for effective chicken production, environmental sustainability and food hygiene.

Keywords: *Prosopis africana*, medicinal plants, phytochemicals, food safety, antibiotics.

***Corresponding Author:** Alagbe Olujimi John, Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India, e-mail: dralagbe@outlook.com

Received: 7 February 2024;

Accepted: 28 March 2024;

Published: 30 April 2024.

1. Introduction

Public concern over potential antibiotic resistance risks related to human health has driven interest in poultry nutrition and adoption of antibiotic free feeding systems (Ulrich & Arne, 2022). This has led to the development of feed additives such as *Prosopis africana* extracts that can be used as in-feed antibiotic alternatives in livestock feeding strategies (Lius, 2022; Jan, 2021). The extracts have a wide range of potential benefits all targeting the enhancement of performance of livestock.

The medium-sized, annual savannah tree Prosopis africana is a member of the fabaceae family. The majority of Asia and Africa (both East and West Africa) are home to the plant (Kolapo et al., 2009). There are roughly forty-five species of it and it can reach heights of 4 to 20 meters. It has a thick tap root that expands quickly and deeply into the earth (Orwa et al., 2009). The plant's leaves range in size from 12 mm to 30 mm and are bipinnate, with 9–16 oblong leaf pairs (Aremu et al., 2007). When dried, the seeds of Prosopis africana can open freely inside a yellow-intermeshed pod (Weber et al., 2008). According to Kolapo et al. (2009), the plants' seeds, leaves, roots and stem

How to cite (APA):

Alagbe, O.J. (2024). *Prosopis africana* extracts as potential natural alternatives to synthetic antibiotics and a key for sustainable broiler production: A review. *Research in Agricultural & Veterinary Sciences*, 8(1), 5-12
<https://doi.org/10.62476/ravs8105>

bark have lots in phytochemicals and are typically used to treat bronchitis, fever, gonorrhea, toothache, stomachache, dysentery and body aches.

The variances in plant extract effectiveness can be attributed to a variety of factors, including climate, location, harvest stage and storage conditions, as well as variations in the chemical content of the extracts made from *Prosopis africana* (Jan, 2020). Numerous phytochemicals found in *P. africana* extracts have been shown to have a variety of pharmacological effects, including anti-inflammatory, anti-helminthic, antioxidant, antimicrobial, immune-stimulatory, anti-fungal and hepatoprotective qualities (Kalemba & Kunicka, 2003).

Essential minerals like potassium, phosphorus, calcium, manganese, magnesium, zinc and copper are present in *Prosopis africana* (Ajiboye *et al.*, 2013). According to Yanwuyi *et al.* (2010), these minerals are critical for the activation of essential enzymes and for protecting against a state of oxidative stress. Additionally, *P. africana* contains the amino acids methionine, lysine, threonine, leucine, alanine, phenyl alanine, cysteine and proline, as well as vitamins A, B2, B5, B6, B9 and B12. These nutrients are essential for reducing cellular damage, scavenging free radicals, promoting collagen formation and further enhancing antioxidant activities (Alagbe, 2022).

Considering the wealth of potential found in *Prosopis africana* and the growing awareness of the need to offer natural antibiotic substitutes instead of manufactured ones. This review summarizes earlier research and emphasizes the need of advancing food safety and sustainable livestock practices.

2. The phytochemical makeup of extracts from *Prosopis africana* phenols

Due to their antioxidant and free-radical scavenging qualities, phenolic compounds help keep animals healthy (Dai & Mumper, 2010). Phenols work by depriving bacteria of their substrate, which causes fragmentation of the bacterial cytoplasm (Cowan 1999). In their investigations, Cushnie and Lamb (2005) and Davies *et al.* (1998) found that the inhibition of enzymes by oxidized substrate was the cause of phenols' potency against bacteria.

Alkaloids

According to Nath *et al.* (2018) and Singh (2008), alkaloids are naturally occurring organic molecules that have a nitrogen heteroatom in their framework. According to Ayushi *et al.* (2021) and Wink (2016), alkaloids may have a variety of medicinal effects on birds, including antibacterial, analgesic and antioxidant qualities. It has been found that alkaloids prevent the synthesis of proteins and RNA (Nath *et al.*, 2018).

Flavonoids

According to Cowan (1999), the carbonyl group of flavonoids has hydroxyl groups at positions 3 and 7. Moreover, it has been proposed that they possess antioxidant and antibacterial qualities. Additionally, they may attach to bacterial cell proteins, inhibiting and deactivating enzymes (Cushnie & Lamb 2005).

Tannins

Every portion of a plant, including the stem, bark, roots, leaves, fruits and flowers, contains tannins (Lim *et al.*, 2006). The capacity of tannins to form complexes with proteins and polysaccharides via covalent and hydrogen bonding is one of their primary

characteristics (Szallasi, 2005). Enzyme inactivation results from tannins' capacity to bind to surface proteins and form complexes that do so.

Enzyme inactivation results from tannins' capacity to bind to surface proteins and form groups that do so. According to Lim et al. (2006), it also causes membrane rupture and substrate restriction as a result of complex formation.

Saponins

A class of phytochemicals known as saponins is one of the main defense mechanisms plants have against microbial, fungal and insect invasion. Whilst the quantity and concentration of saponins in plants differ from species to species, they are present in the majority of plant species (González-Lamothe *et al.*, 2009). According to Morrissey and Osbourn (1999), saponins function by building compounds with sterols or polysaccharides inside the microbial cell membrane, which destroys the integrity of the cytoplasmic membrane.

3. Pharmacological properties of *Prosopis africana*

Antimicrobial properties

A study carried out by Badri et al. (2016) showed essential oils from *Prosopis africana* have the capacity to inhibit the activities of some bacteria's, such as: *Shigella flexneri*, *Salmonella typhi*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Klebsiella pneumonia*, *Enterococcus faecalis*, *Listeria monocytogenes* and *Bacillus cereus* due to the presence of phytochemicals like phenols, tannins and flavonoids which have been suggested to perform multiple biological activities against pathogenic organisms. Another studies have showed that methanolic and ethanolic extracts from the leaves, stem bark and roots of *Prosopis* are effective against *Salmonella typhi*, *Proteus mirabilis*, *Enterococcus faecalis* and *Listeria monocytogenes* (Tajbakhsh *et al.*, 2015).

The properties of antioxidants

Extracts that are aqueous, methanolic and ethanolic have a number of medicinal uses and have the ability to scavenge free radicals that can lead to infections in an animal's body (Darogba *et al.*, 2009). Additionally, according to Saad et al. (2017), prosopis oil demonstrated efficacy against *Erwinia* spp., *C. albicans*, *P. vulgaris*, *E. coli* and *Shigella* spp. Additionally, research has demonstrated that adding 400 mg/kg of prosopis oil to broiler diets can have a major impact on the activity of enzymes such as glutathione peroxidase, catalase, reduced glutathione, superoxide dismutase and malondialdehyde (Alagbe, 2023).

Hypolipidemic characteristics

According to earlier research by Rahman et al. (2011), *Prosopis* extracts administered to rats at doses of up to 600 mg/kg significantly affected their levels of triglycerides, total cholesterol, low density lipoprotein, very low density lipoprotein and atherogenic index (meat safety index), indicating a potential preventive effect against cardiovascular diseases. Additionally, according to Alagbe et al. (2022), adding 800 mg/kg of *Prosopis africana* essential oil to the diet can lower the amount of saturated fat and raise the amount of polyunsaturated fat in broiler meat (Alagbe, 2022).

4. Conclusion

Medicinal plants are infinitely capable of producing phytochemicals, which are chemical substances with a vast range of possible benefits, all aimed at improving bird performance. *Prosopis africana* extracts contain several substances, which have been demonstrated to be safe, efficacious and environmentally benign. Additionally, studies have revealed that *P. africana* extract includes considerable amounts of phenols, tannins, alkaloids, terpenoids and flavonoids, which provide them antibacterial, antioxidant, antifungal, antiviral and other beneficial properties. Additionally, adding nutrition to broiler feed may improve the birds' performance. These will increase the output of chickens and lower the rising number of antibiotic-resistant illnesses.

Table 1. Some bioactive compounds in *Prosopis africana* oil and their properties (Alagbe, 2023)

Active compounds	Medicinal properties	References
Prosogerin A (6 Methoxy-7-hydroxyl dioxyflavone)	Antibacterial, antioxidant and immune-stimulatory	Guimarães (2010)
Prosogerin B (2,4 dihydroxyl -5 methyl dioxychacon)	Antimicrobial, hepato-protective and antioxidant	Jim'enez-Arellanes et al. (2003)
2,4-bis (1-phenylethyl) phenol	Antioxidants, antifungal	Kokoska et al. (2002)
β -phenethylamine	Antifungal and immune-stimulatory	Lin et al. (2000)
2,4,6-tris(1-phenylethyl) phenol	Antibacterial, antiviral	Loizou et al. (2010)
α -pinene	Antioxidant and digestive stimulants	Alagbe et al. (2022)
α -terpinene	Antioxidant, antibacterial,	Alagbe et al. (2023)
β -pinene	Antioxidant, antimicrobial	Egunyomi and Oladunjoye (2012)
β -myrcene	Antioxidant, digestive stimulants	Edeoga et al. (2005); Alagbe (2022)
α -phellandrene	Antioxidant, Antifungal and antimicrobial	Idowu et al. (2006)
α -terpinolene	Antioxidant, antibacterial	Guimarães (2010)
γ -terpinene	Antioxidant, immune-stimulatory	Jim'enez-Arellanes et al. (2003)
1-terpineol	digestive stimulants	Lalitha et al. (2011)
4-terpineol	digestive stimulants	Lin et al. (2000)
Humulene	Antibacterial, anti-helminthic	Loizou et al. (2010)
Caryophyllene	digestive stimulants	Alagbe et al. (2022)
Copaene	Antimicrobial	Wild (1994)
Cis-linaloxide	Antifungal	Xu and Chang (2007)
α -Selinene	Antioxidant, anti-inflammatory	Shai et al. (2008)
γ -Elemene	digestive stimulants	Shaheen et al. (2005)
α -Gurjunene	Immuno-stimulatory, antioxidant	Özçelik et al. (2011)
β -Elemene	Antimicrobial, antioxidant, digestive stimulants	Singh et al. (2023); Omokore and Alagbe (2019)
β -Cyclocitral	Anti-inflammatory, digestive stimulants	Kolapo et al. (2009)
3-hexenyl-2-methylbutanoate	Anti-bacterial	Inngjerdingen et al. (2004)
Exo-methyl-camphenilol	Antimicrobial	Ezike et al. (2010)
Caryophyllene oxide	Antibacterial, antioxidant	Enright et al. (2002)
Benzene -1-methoxy-2-methyl	Antifungal	Fabricant and Farnsworth (2001)
Napthalene, 1,2 hydro-1,1, 6 – trimethyl	Antimicrobial	Inngjerdingen et al. (2004)
Cycloheptasiloxane, tetradecamethyl	Anti-inflammatory	Chua (2013); Daniel et al. (2023)
Nerolidyl acetate	Anti-inflammatory	Azmir et al. (2013)
1-Cyclohexene-1- butanal, alpha, 2, 6, 6- tetramethyl	Anti-inflammatory	Atanasov et al., 2015; Alagbe (2024)

Table 2. Effects of *Prosopis africana* extracts on the general performance of birds

Type of extracts used	Concentration used	Effect on broilers
<i>Prosopis africana</i> essential oil	400 mg/kg	Increased body weight gain
		Increased pancreatic enzyme production
		Improved nutrient utilization
		Increased feed intake
		Reduced mortality rate
		Increased in carcass weight
<i>Prosopis africana</i> essential oil	800 mg/kg	Increased production of polyunsaturated fatty acid in meat sample
		Improved sensory attributes of meat
		Increased in pack cell volume, red blood cell, haemoglobin, white blood cell, total protein, amongst others
		Scavenging free radicals
		Increasing antibody titres in birds after vaccination
		Decrease in population of pathogenic microorganisms
<i>Prosopis africana</i> stem bark extracts (aqueous)	8ml/litre of water	Increased final body weight gain, feed intake and <i>Lactobacillus sp</i> count in the gut of broilers

References

- Ajiboye, A.A., Agboola, D.A., Fadimu, O.Y. & Afolabi, A.O. (2013). Antibacterial, phytochemical and proximate analysis of *Prosopis africana* (Linn) seed and pod extract. *FUTA Journal of Research in Sciences*, 1, 101-109.
- Alagbe, J.O. (2021). *Prosopis africana* stem bark as an alternative to antibiotic feed additives in broiler chicks diets: Performance and Carcass characteristics. *Journal of Multidimensional Research and Reviews*, 2(1), 64-77.
- Alagbe, J.O. (2023). Investigating the effects of dietary supplementation of eucalyptus camaldulensis essential oil (eceo) on the growth performance, nutrient digestibility and caecal fermentation of weaned rabbits. *Research in: Agricultural & Veterinary Sciences*, 7(3), 139 - 148.
- Alagbe, J.O. (2023). Investigating the effects of dietary supplementation of *Eucalyptus camaldulensis* essential oil on haemato-biochemical indices, immune response and oxidative stress of weaned rabbits. *International Journal of Agriculture and Animal Production*, 4(1), 34 - 46.
- Alagbe, J.O. (2022). *Prosopis africana* (African mesquite) oil as an alternative to antibiotic feed additives on broiler chicken's diets: Haematology and serum biochemical indices. *Central Asian Journal of Theoretical and Applied Sciences*, 3(2), 19-29.
- Alagbe, J.O. (2022). *Prosopis africana* (African mesquite) oil as an alternative to antibiotic feed additives on broiler chickens diets: Performance and nutrient retention. *Discovery*, 58(314), 134 -142.
- Alagbe, J.O. (2022). Use of medicinal plants as a panacea to poultry production and food security: A review. *American Journal of Technology and Applied Sciences*, 1, 24-36.

- Alagbe, J.O., Anorue, D.N., Shittu, M.D., Ramalan, S.M., Faniyi, T.O. & Ajagbe, A.D. (2024). Growth performance and physiological response of weaned pigs fed diet supplemented with novel a phytogenics. *Brazilian Journal of Science*, 3(1), 43-57.
- Alagbe, J.O., Omokore, E.A. (2019). Effect of replacing soya bean meal with Indigofera zollingeriana leaf meal on the performance and carcass characteristics of growing rabbits. *International Journal of Multidisciplinary Research and Development*, 6(5), 74-77.
- Alagbe, J.O. (2024). Novel phytogenics' impact on weaned pigs' growth performance, heamatology and serum biochemical indicators. *Black Sea Journal of Agriculture*, 7(2), 1-2.
- Alagbe, O.J. (2023). Sensory evaluation and fatty acid composition of broiler chickens fed diets containing with Prosopis africana essential oil. *Journal of Healthcare and Biomedical Science*, 1(2), 35-45.
- Alagbe, O.J., Daniel, N.A., Shittu, M.D., Emiola, A., Akande, T. & Adegoke, A.E. (2023). Impact of dietary supplementation of Carica papaya essential oil on the blood chemistry of broiler chickens. *Science Letters*, 11(3), 105 - 110.
- Atanasov, A.G., Waltenberger, B., Pferschy-Wenzig, E.M., Linder, T., Wawrosch, C., Uhrin, P. & Stuppner, H. (2015). Discovery and resupply of pharmacologically active plant-derived natural products: A review. *Biotechnology Advances*, 33(8), 1582-1614.
- Atawodi, S.E., Ogunbusola, F. (2009). Evaluation of anti-trypanosomal properties of four extracts of leaves, stem and root barks of Prosopis africana in laboratory animals. *Biokemistri*, 21(2).
- Ayanwuyi, L.O., Yaro, A.H. & Abodunde, O.M. (2010). Analgesic and anti-inflammatory effects of the methanol stem bark extract of Prosopis africana. *Pharmaceutical Biology*, 48(3), 296-299.
- Azmir, J., Zaidul, I.S.M., Rahman, M.M., Sharif, K.M., Mohamed, A., Sahena, F. & Omar, A.K.M. (2013). Techniques for extraction of bioactive compounds from plant materials: A review. *Journal of Food Engineering*, 117(4), 426-436.
- Chua, L.S. (2013). A review on plant-based rutin extraction methods and its pharmacological activities. *Journal of Ethnopharmacology*, 150(3), 805-817.
- Cowan, M.M. (1999). Plant products as antimicrobial agents. *Clinical Microbiology Reviews*, 12(4), 564-582.
- Cushnie, T.T., Lamb, A.J. (2005). Antimicrobial activity of flavonoids. *International Journal of Antimicrobial Agents*, 26(5), 343-356.
- Dai, J., Mumper, R.J. (2010). Plant phenolics: Extraction, analysis and their antioxidant and anticancer properties. *Molecules*, 15(10), 7313-7352.
- Davies, D.G., Parsek, M.R., Pearson, J.P., Iglewski, B.H., Costerton, J.W. & Greenberg, E.P. (1998). The involvement of cell-to-cell signals in the development of a bacterial biofilm. *Science*, 280(5361), 295-298.
- Edeoga, H.O., Okwu, D.E. & Mbaebie, B.O. (2005). Phytochemical constituents of some Nigerian medicinal plants. *African Journal of Biotechnology*, 4(7), 685-688.
- Egunyomi, A.S. Oladunjoye, S. (2012). Studies on the chemical composition and Nutritive value of the fruit of African Star Apple. *African Journal of Agricultural Research*, 7(31), 4256-4258.
- Enright, M.C., Robinson, D.A., Randle, G., Feil, E.J., Grundmann, H. & Spratt, B.G. (2002). The evolutionary history of methicillin-resistant Staphylococcus aureus (MRSA). *Proceedings of the National Academy of Sciences*, 99(11), 7687-7692.
- Ezike, A.C., Akah, P.A., Okoli, C.O., Udegbunam, S., Okwume, N., Okeke, C. & Iloani, O. (2010). Medicinal plants used in wound care: A study of Prosopis africana (Fabaceae) stem bark. *Indian Journal of Pharmaceutical Sciences*, 72(3), 334-339.
- Fabricant, D.S., Farnsworth, N.R. (2001). The value of plants used in traditional medicine for drug discovery. *Environmental Health Perspectives*, 109(suppl 1), 69-75.

- Guimarães, L. (2010). Óleos essenciais de lippia sidoides Cham., alomia fastigata (Gardner) Benth, ocotea odorífera (vell.) Rower, mikania glauca Mart. E cordia verbenaceae DC: Identificação e quantificação química, caracterização das estruturas secretoras, atividades antioxidante e antibacteriana. 2010. 225 f (Doctoral dissertation, Tese (Doutorado em Agroquímica)-Universidade Federal de Lavras, Lavras).
- Idowu, T.O., Iwalewa, E.O., Aderogba, M.A., Akinpelu, B.A. & Ogundaini, A.O. (2006). Antinociceptive, anti-inflammatory and antioxidant activities of eleagnine: An alkaloid isolated from Chrysophyllum albidum seed cotyledons. *Journal of Biological Science*, 6, 1029–1034.
- Inngjerdingen, K., Nergård, C.S., Diallo, D., Mounkoro, P.P. & Paulsen, B.S. (2004). An ethnopharmacological survey of plants used for wound healing in Dogonland, Mali, West Africa. *Journal of Ethnopharmacology*, 92(2-3), 233-244.
- Jimenez-Arellanes, A., Meckes, M., Ramirez, R., Torres, J. & Luna-Herrera, J. (2003). Activity against multidrug-resistant Mycobacterium tuberculosis in Mexican plants used to treat respiratory diseases. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*, 17(8), 903-908.
- Kokoska, L., Polesny, Z., Rada, V., Nepovim, A. & Vanek, T. (2002). Screening of some Siberian medicinal plants for antimicrobial activity. *Journal of Ethnopharmacology*, 82(1), 51-53.
- Kolapo, A.L., Okunade, M.B., Adejumbi, J.A. & Ogundiya, M.O. (2009). Phytochemical composition and antimicrobial activity of Prosopis africana against some selected oral pathogens. *World Journal of Agricultural Sciences*, 5(1), 90-93.
- Lalitha, V., Kiran, B. & Raveesha, K.A. (2011). Antifungal and antibacterial potentiality of six essential oils extracted from plant source. *Int. J. Eng. Sci. Technol*, 3(4), 3029-3038.
- Lin, F.Y., Vo, A.H., Phan, V.B., Nguyen, T.T., Bryla, D., Tran, C.T. & Robbins, J.B. (2000). The epidemiology of typhoid fever in the Dong Thap Province, Mekong Delta region of Vietnam. *The American Journal of Tropical Medicine and Hygiene*, 62(5), 644-648.
- Loizou, S., Lekakis, I., Chrousos, G.P., & Moutsatsou, P. (2010). β -Sitosterol exhibits anti-inflammatory activity in human aortic endothelial cells. *Molecular Nutrition & Food Research*, 54(4), 551-558.
- Nnadozie Anorue, D., Ubong, F. & John, A.O. (2023). Investigating the effect of pawpaw (Carica papaya) essential oil dietary supplementation on the growth performance and carcass characteristics of broilers. *Research in: Agricultural & Veterinary Sciences*, 7(3), 164-174.
- Özçelik, B., Kartal, M. & Orhan, I. (2011). Cytotoxicity, antiviral and antimicrobial activities of alkaloids, flavonoids and phenolic acids. *Pharmaceutical Biology*, 49(4), 396-402.
- Sandasi, M., Leonard, C.M. & Viljoen, A.M. (2010). The in vitro antibiofilm activity of selected culinary herbs and medicinal plants against Listeria monocytogenes. *Letters in Applied Microbiology*, 50(1), 30-35.
- Shaheen, A.Y., Sheikh, A.A., Rabbani, M., Aslam, A., Bibi, T., Liaqat, F. & Rehmani, S.F. (2015). Antibacterial activity of herbal extracts against multi-drug resistant Escherichia coli recovered from retail chicken meat. *Pakistan Journal of Pharmaceutical Sciences*, 28(4), 1295-1300.
- Shai, L.J., McGaw, L.J., Masoko, P. & Eloff, J.N. (2008). Antifungal and antibacterial activity of seven traditionally used South African plant species active against Candida albicans. *South African Journal of Botany*, 74(4), 677-684.
- Sharma, S., John, A.O., Xing, L., Ram, S. & Amita, K. (2022). Comparative analysis of ethanolic Juniperus thurifera leaf, stem bark and root extract using gas chromatography and mass spectroemetry. *International Journal of Agriculture and Animal Production*, 2(6), 18-27.

- Singh, J. (2008). Maceration, percolation and infusion techniques for the extraction of medicinal and aromatic plants. *Extraction Technologies for Medicinal and Aromatic Plants*, 67, 32-35.
- Wild, R. (1994). The complete book of natural and medicinal cures. *Complement Health Pract Rev*, 3, 139.
- Xu, B.J., Chang, S.K. (2007). A comparative study on phenolic profiles and antioxidant activities of legumes as affected by extraction solvents. *Journal of Food Science*, 72(2), S159-S166.